

## Study of the Ground-state Configuration of Neutron-rich Aluminium Isotopes through Coulomb Breakup

*S. Chakraborty<sup>1</sup>, U. Datta Pramanik<sup>1#</sup>, T. Aumann<sup>2,3</sup>, G. De Angelis<sup>4</sup>, S. Beceiro<sup>5</sup>, T. Le Bleis<sup>2</sup>, K. Boretzky<sup>2</sup>, D. Cotrina-Gil<sup>5</sup>, C. Caesar<sup>2</sup>, M. Chariter<sup>6</sup>, W. N. Catford<sup>7</sup>, D. Gonzalez-Diaz<sup>2</sup>, H. Emiling<sup>2</sup>, P. Diaz-Fernandez<sup>5</sup>, L. M. Fraile<sup>9</sup>, O. Ershova<sup>2</sup>, H. Geissel<sup>2</sup>, M. Heil<sup>2</sup>, M. Heine<sup>2</sup>, B. Jonson<sup>10</sup>, A. Kelic<sup>2</sup>, H. T. Johansson<sup>10</sup>, R. Krucken<sup>11</sup>, T. Kroll<sup>3</sup>, C. Langer<sup>2</sup>, Y. Leifell<sup>2</sup>, G. Munzenberg<sup>2</sup>, J. Marganec<sup>15</sup>, C. Nociforo<sup>2</sup>, M. A. Najafi<sup>12</sup>, V. Panin<sup>2</sup>, R. Plag<sup>2</sup>, H. A. Pol<sup>5</sup>, S. Paschalis<sup>13</sup>, A. Rahman<sup>1</sup>, R. Reifarh<sup>2</sup>, D. Rossi<sup>2</sup>, H. Simon<sup>2</sup>, C. Scheidenberger<sup>2</sup>, S. Typel<sup>2</sup>, J. Taylor<sup>6</sup>, Y. Togano<sup>14</sup>, V. Volkov<sup>2</sup>, H. Weick<sup>2</sup>, A. Wagner<sup>8</sup>, F. Wamers<sup>2</sup>, J. Winfield<sup>2</sup>, M. Zoric<sup>2</sup> and the S306 collaboration.*

<sup>1</sup>Saha Institute of Nuclear Physics, Kolkata, India; <sup>2</sup>GSI, Darmstadt, Germany; <sup>3</sup>TU, Darmstadt; <sup>4</sup>INFN, Legnaro, Italy; <sup>5</sup>Univ. of Santiago de Compostela, Spain; <sup>6</sup>Germany; <sup>7</sup>University of Liverpool, UK; <sup>8</sup>HZDR Dresden-Rossendorf, Germany; <sup>9</sup>UCM, Madrid, Spain; <sup>10</sup>Chalmers University of Technology, Sweden; <sup>11</sup>TU, Munich, Germany; <sup>12</sup>KVI Groningen, Netherland; <sup>13</sup>LBNL, Berkeley, USA; <sup>14</sup>RIKEN, Japan; <sup>15</sup>EMMI, GSI, Darmstadt, Germany;

In 1949 Mayer and Jensen successfully explained the magic number in the nuclei using the spin-orbit interaction. This theory explained a vast amount of reachable experimental data in the next three decades until the ground breaking experiment by Thibault et. al.[1] in 1975 on neutron rich nuclei near the neutron magic number  $N \sim 20$ . In this region of nuclei, the ground states are dominated by intruder configurations from upper  $pf$  orbits [2]. Hence, this region of nuclear chart is called “*island of inversion*”. Nuclei like  $^{34,35}\text{Al}$  are lying close to this *island of inversion*. Little experimental information on ground state configuration of those isotopes are available in literature [3,4] regarding their intruder  $pf$ -shell contribution. Coulomb excitation is a direct probe for studying the ground state configuration of loosely bound nuclei [5,6]. Experiment S306 was performed using the existing RIB facility at GSI, Darmstadt to study the properties of the nuclei in and around the  $N \sim 20$  *island of inversion* through electromagnetic excitation. Short-lived radioactive nuclei were produced by the fragmentation of  $^{40}\text{Ar}$  beam (at 531 MeV/u) on Be (8 gm/cm<sup>2</sup>) production target at fragment separator (FRS). Secondary beam from FRS, containing  $^{34,35}\text{Al}$  were allowed to fall on various targets [Pb for electromagnetic excitation, Carbon for nuclear excitation and without target for reactions induced by detector materials] at Cave C with the exclusive set-up for kinematically complete measurement, the FRS -LAND set-up.

The incoming beam was identified uniquely by energy loss and ToF measurements before the reaction target along with the known magnetic rigidities of FRS. Neutrons and  $\gamma$ -rays from the de-exciting projectile or projectile like fragments were detected by the LAND and the  $4\pi$ -Crystal Ball spectrometer, respectively. Reaction fragments were tracked via the Silicon Strip Trackers and GFI detectors placed before and after the magnetic spec-

trometer (ALADIN), respectively. Finally, mass of




Fig. 1:  $\gamma$ -sum spectra of  $^{33}\text{Al}$  obtained after Coulomb breakup of  $^{34}\text{Al}$  in lead target.

the outgoing fragments were identified by reconstructing the magnetic rigidities inside ALADIN event by event using the tracker programme developed at SINP. Fig. 1 shows the  $\gamma$ -sum spectra of  $^{33}\text{Al}$  obtained after Coulomb breakup of  $^{34}\text{Al}$ . The spectrum was obtained in coincidence with  $^{33}\text{Al}$  fragments and one neutron. These observed  $\gamma$ -lines are in agreement with the characteristic  $\gamma$ -rays of  $^{33}\text{Al}$  reported in the literature [7]. Comparison of the measured Coulomb breakup differential cross-sections with the theoretically calculated cross-sections will help us to pin down the ground state configuration and shell inversion in these neutron-rich nuclei.

### References

- [1] C. Thibault, et al., Phys. Rev. C 12 (1975) 644.
- [2] T. Motobayashi, et al., Phys. Lett. B 346 (1995) 9.
- [3] P. Himpe, et al., Phys. Lett. B 643 (2006) 257.
- [4] C. Nociforo, et. al. Phys. Rev. C 85 (2012) 044312.
- [5] T. Nakamura, et. al. Phys. Rev. Lett. 83, (1999) 1112.
- [6] U. DattaPramanik, et al., Phys. Lett. B 551 (2003) 63.
- [7] [www.nndc.bnl.gov/](http://www.nndc.bnl.gov/)

\*Supported by Alexander von Humboldt (AvH) foundation.

#ushasi.dattapramanik@saha.ac.in